

Original Research Paper

# Diabetic Retinopathy: Changes in Levels of Orosomucoids in Patients Supplemented with Beta Glucan and Vitamin D

<sup>1</sup>Richter Josef, <sup>2</sup>Závorková Martina, <sup>1</sup>Stiborova Ivana, <sup>1</sup>Král Vlastimil and <sup>3</sup>Vetvicka Vaclav

<sup>1</sup>Zdravotní ústav se sídlem v Ústí nad Labem, Ústí nad Labem, Czech Republic

<sup>2</sup>Oční klinika UJEP Masarykova nemocnice, Krajská zdravotní, a.s., Ústí nad Labem, Czech Republic

<sup>3</sup>University of Louisville, Department of Pathology, Louisville, KY, USA

## Article history

Received: 12-03-2019

Revised: 27-03-2019

Accepted: 13-04-2019

Corresponding Author:

Vaclav Vetvicka

University of Louisville,

Department of Pathology, 511

South Floyd Street, Louisville,

KY 40202, USA

Email: vaclav.vetvicka@louisville.edu

**Abstract:** The role of beta glucan in stimulating of various biological reactions is getting more and more attention. In this report, we focused on the effects of oral supplementation with beta glucan and vitamin D on changes in serum levels of orosomucoids in patients with diabetic retinopathy. We measured the level of orosomucoids and evaluated the effects of oral supplementation. In this study we report significant decrease of orosomucoids levels after the end of the three-month treatment. Our results support the fact that continual supplementation with glucan is necessary for diabetes melitus and diabetic retinopathy patients.

**Keywords:** Glucan, Diabetic Retinopathy, Vitamin D, Treatment, Orosomucoids

## Introduction

Prevalence of Diabetes Mellitus (DM) shows steep incline, both in DM Type 1 (DM1) and less steeply in DM Type 2 (DM2) (Duh *et al.*, 2017; Muhammad *et al.*, 2017). Diabetic patients are endangered by various complications, influencing the quality of their life functions. Diabetic Retinopathy (DR) holds the most common complications of DM with current prevalence reaching more than 10 million worldwide (El-Beblawy *et al.*, 2016). Regulation of immune functions and control of inflammatory processes are decisive in maintaining normal retinal functions. Chronic parainflammatory process leads to initiation and manifestation of several impairments, including DR (Nita *et al.*, 2014). Factors directly related to prevention of DM and reduction of incidence are: (1) quality of lifestyle, (2) nutrition, (3) reduction of obesity, (4) related metabolic syndrome. Inducing production of several proteins, gradually occurring by weight increase, is dependent on fat cells. It is increased by oxidative stress leading to dysregulation of normal parainflammatory reaction and chronic inflammation (Duh *et al.*, 2017; Hajer *et al.*, 2008; Tsuboi *et al.*, 2018). Recent studies monitoring relations between Orosomucoids (ORM) and pathogenesis of DM found that high levels of ORM are related to regulation of glucose levels. ORM, also known as  $\alpha$ 1-acid

glycoprotein, is one of the acute-phase proteins exhibiting a variety of activities both *in vitro* and *in vivo*.

Risk of DM development increases with weight gain, together with an increase of clinical problems related to metabolic syndrome (Richter *et al.*, 2018a; 2018b; Zhou *et al.*, 2017). Authors emphasize that for evaluation of this relation, it is better to replace the BMI criterium by other measurements (such as WHR) (Zhou *et al.*, 2017; Zavorokova *et al.*, 2018). Incidence of additional comorbidities connected with high weight is also related to significant changes of nonspecific and specific immune responses. These changes involve, among others, Toll-like receptors that are present not only on immunocytes, but also on adipocytes and intestinal cells. They play a part in development of complex balance influencing normal immune system and metabolism (Wolowczuk *et al.*, 2008). Fat cell dysfunction in obese population is one cause of DM2 development (Hajer *et al.*, 2008). Weight reduction and increased physical activity are effective interventions leading to fat cell function improvement.

Several studies described positive correlation between fat weight and levels of proteins signaling inflammation (Engstrom *et al.*, 2003). Weight reduction in obese individuals is closely correlated to decrease elevated levels of inflammation and inflammation-signaling proteins. Proinflammatory cytokines,

produced by adipocytes and preadipocytes (such as TNF- $\alpha$  and IL-6) significantly influence production of these inflammatory proteins: (1) ORM, (2) fibrinogen, (3) alpha 1 antitrypsin, (4) haptoglobin, (5) ceruloplasmin CRP, (6) SAA. These proteins subsequently support the occurrence of numerous diseases (Jensen and Whitehead, 1998). Literature shows that persistent increased levels of acute inflammation proteins can predict significant increase in weight in middle age population. This relation can lead to the assumption about possible subsequent connection with metabolic syndrome, cardiovascular diseases, diabetes and other health problems (Muhammad *et al.*, 2017; Engstrom *et al.*, 2003; Jensen and Whitehead, 1998). Association of elevated levels of ORM, haptoglobin and CRP with an elevated risk of DM development was already described (Duh *et al.*, 2017; Muhammad *et al.*, 2017). This high risk of diabetes related to high levels of acute inflammatory protein can be accepted with one condition: additional possible mechanisms involved must be evaluated and described (Muhammad *et al.*, 2017). High ORM levels on DM1 reflect endothelial dysfunction and subclinical atherosclerosis and can be considered as an early marker of renal injury (El-Beblawy *et al.*, 2016). ORM can regulate food intake and homeostasis of energy (Sun *et al.*, 2016). Modulation of ORM expression can be considered as a new approach in regulation of obesity and subsequent diseases related to metabolic syndrome (Cloetens *et al.*, 2012). Immunomodulatory mechanisms, drug transport, capillar quality maintenance and sphingolipid metabolism are mediated via ORM binding to CCR-5 or to beta chain of hemoglobin (Muhammad *et al.*, 2017; Qin *et al.*, 2016). Levels of ORM correlate with BMI, fat deposits, leptin levels and fasting glucose levels. This suggests that ORM can participate in regulation of energetic balance (Muhammad *et al.*, 2017; Lei *et al.*, 2016). These findings were further confirmed by using mice lacking ORM1. These animals clearly displayed aberrant energy homeostatis including increased body weight and fat mass (Sun *et al.*, 2016).

ORM has several additional modulatory activities: (1) immunomodulation, (2) sphingolipid metabolism, (3) drug transport, (4) keeping of capillar barrier function (Luo *et al.*, 2015). All these findings have resulted in current intensive research in evaluation of ORM functions, particularly in relation to physiology of energy. It is well known that physical stress leads to elevated levels of ORM and subsequent reduction of fatigue upon binding to C-C chemokine receptor type 5 (Lei *et al.*, 2016; Straczkowski *et al.*, 2018). Together with additional proteins of inflammatory response, ORM is a good risk indicator of cardiovascular damage and cardiovascular disease (Lei *et al.*, 2016). ORM is considered to be proof of association between obesity

and periodontitis (Range *et al.*, 2013). Interesting studies measuring ORM in urine of patients with DM1 might be used in induction risk evaluation with urinary tract problems concerning DR patients (Christiansen *et al.*, 2010; Kustan *et al.*, 2016; Talks *et al.*, 2018). Lately, increased attention is focused on evaluation of ORM I and II subclasses. Fascinating studies describing increased ORM I expression in odontogenic myxoma and particularly regulatory abilities ORM II in hepatocellular carcinome (Garcia-Munoz *et al.*, 2012) inspired the possibility to assess levels of ORM I and ORM II during diabetes. This is not only in relation of possible preventive actions, but also in monitoring of possible risk of complications known during DM.

Several studies demonstrated associations between levels of vitamin D and increased risk of development of DM2 (Lu *et al.*, 2018; Spiro and Buttriss, 2014). Vitamin D supplementation, resulting in changes of its levels, results in significant decrease of risk of DM2 (Zavorkova *et al.*, 2018; Luo *et al.*, 2015). Numerous studies found deficits in vitamin D levels not only in European countries, but worldwide (Tessari and Lante, 2017). Some of these countries already established a program of food fortification with vitamin D towards improved quality of health of entire populations ((Tessari and Lante, 2017). A complex preventive step aiming on reduction of risks of DM2 development is using supplementation with vitamin D, C and E (Garcia-Bailo *et al.*, 2011). In addition, the importance of affecting inflammatory reaction as DM2 induction prevention, is proposed (Garcia-Bailo *et al.*, 2011; Bashir and Choi, 2017).

Another possibility leading to weight reduction is influencing inflammatory response by addition of beta glucan, resulting in reduction of inflammatory response mediated by preadipocytes and adipocytes. The fact that a diet fortified with fiber containing high glucan content increased metabolic readiness and can be used in DM2 patients is well established (Nita *et al.*, 2014; McFarlin *et al.*, 2013).

Some studies describe effects of food supplemented with glucan and psyllia on DM2 occurrence (Aoe *et al.*, 2017; McRae, 2018). Beta glucan is established to be an important part of nutrition, which is able to modulate dysregulation of metabolism connected with metabolic syndrome (El Khoury *et al.*, 2012; Lyon and Reichert, 2010; Aoe *et al.*, 2017). In addition, glucan can significantly decrease food consumption, regulate glycemic index of food and positively affect glucose metabolism in patients with DM2 (Straczkowski *et al.*, 2018; Bashir and Choi, 2017; McRae, 2018). In our group of patients with DM2 and DR we demonstrated significant changes of lipid metabolism caused by improvements of vitamin D levels and after glucan application (Zavorkova *et al.*, 2018). In

the current study we follow-up the previous work and focus on evaluation of ORM levels after glucan and vitamin D supplementation.

## Materials and Methods

### Protocol

We explained the experimental protocol and obtained consent forms from all participating patients. This study was Institutional Review Board (Regional Masaryk University) approved and performed in full agreement with the Helsinki declaration (revised version 2000.09.01) and in full compliance with the Czech Republic's clinical testing rules.

### Patients

Our initial group of patients consisted of 52 patients with Diabetic Retinopathy (DR) on one or both eyes. In addition, this disease was further complicated by Diabetic Macular Edema (DME). Initial criteria included Central Thickness of Retina (CTR) over 250  $\mu\text{m}$  and best corrective eye sharpness 4/6.3 to 4/40. Excluding criteria were classic laser photocoagulation of macula less than 3 months before the start of the study, previous intraocular treatment via anti-VEGF antibodies less than 6 months before the start of the study, additional macular disease and situation after pars plana vitrectomy. Average CTR at the start of the study was  $448.2 \pm 132.3$   $\mu\text{m}$  (median 441.5), average best corrective eye sharpness was  $63.1 \pm 13.4$  (median 65) of letters.

A group of a total of 52 patients diagnosed with diabetic retinopathy was divided into three groups. Group A (20 patients) was supplemented with glucan and vitamin D, Group B (20 patients) was supplemented with vitamin D and placebo and Group C (12 patients) was a control group. In all patients we took blood at regular intervals for a 3 month period. All immunological and biochemical observations necessary to follow the development of disease, together with basic treatment using subthreshold micropulse laser of retina by an eye doctor was performed.

At the beginning of the study, the average BMI was  $31.34 \pm 5.1$ , ABSI levels  $0.0856 \pm 0.00076$ . Only 4 patients (7.24%) had an optimal weight, 33 patients (60%) has BMI over 30 and the rest had BMI over 32.5 (32.76%). The starting level of vitamin D was  $14.19 \pm 5.8$  ng/mL. Three months of supplementation resulted in an increase to  $22.45 \pm 7.46$  ng/ml ( $p = 0,001$ ), additional 3 months increased the vitamin D levels to  $29.21 \pm 14.16$  ng/ml, which almost reached recommended levels. After six months, 3 patients had vitamin D levels below 10 ng/ml, 19 patients had vitamin D levels over 30 ng/ml.

### Glucan

Yeast-derived insoluble Glucan #300 (>85% dry w/w basis) was purchased from Transfer Point (Columbia, SC, USA). This glucan contains 96% carbohydrates and 2.1% proteins. Neutral sugar analysis confirmed 91.3% glucose and 8% mannose. 500 mg of glucan was taken on an empty stomach, followed by 100 mL of water and a 30-minute rest prior to any food intake. The interval of this food supplementation was 3 months. Placebo was identical in size, shape and color.

### Vitamin D

Vitamin D (colecaciferol, D3) was manufactured by Merck (Darmstadt, Germany). One ml of solution contains 20,000 IU of vitamin D3, one drop contains 500 IU. All patients were instructed to ingest vitamin D with fat-containing food. The dose used in our experiments was based on weight, sex, age and phototype with 80% reduction during summer (May to September).

### Testing


Serum levels of ORM were tested using diagnostic kits and controls manufactured by Siemens (Siemens Health Care Diagnostics, Newark, DE, USA). All measurements were performed immediately after blood collection.

### Statistical Analysis


Paired *t* test statistical significance was evaluated (GraphPad Prism 5.04; GraphPad Software, USA). An average and standard deviation was evaluated after determining standard value composition (D'Agostino, Pearson). In case of nonstandard composition, values were converted into logarithms.

## Results

Our findings of levels of ORM levels in serum are shown in Fig. 1. Average levels at the beginning of the study were equal in all tested groups. Group A had an average level of ORM  $97.6 \pm 23.86$  g/L, Group B  $100.41 \pm 30.91$  g/L and Group C  $99.45 \pm 24.8$  g/L. Our findings at 3 and 6 months were different. In Group A we found significantly different values ( $P = 0.001$ ) at month 3. Ending glucan supplementation with subsequent vitamin D supplementation resulted in slow increase of ORM levels to original levels of  $102.21 \pm 26.96$  g/L ( $p = 0.0199$ ). Our findings in Group B supplemented with vitamin D and placebo were identical throughout the entire study –  $100.41 \pm 30.91$  g/L at the beginning,  $99.6 \pm 23.97$  g/L after 3 months and  $96.73 \pm 21.05$  g/L after 6 months. Similarly, levels of ORM in Group C were not changed –  $99.45 \pm 24.0$  g/L,  $95.9 \pm 24.0$  g/L and  $95.0 \pm 25.0$  g/L, respectively. Differences between levels found in Groups B and C were tested to be statistically insignificant.



**Fig. 1:** Effects of glucan and vitamin D supplementation on levels of ORM. Group A – glucan and vitamin D; Group B – placebo and vitamin D; Group C – control. Average levels at the beginning of the study were equal in all tested groups. Our findings at 3 and 6 months were different. In Group A we found significantly different values at month 3. Ending glucan supplementation with subsequent vitamin D supplementation resulted in slow increase of ORM levels to original levels. Findings in Group B supplemented with vitamin D and placebo and in Group C were identical throughout the entire study



**Fig. 2:** BMI in tested groups. GLU – glucan and vitamin D (at the beginning and the end of the study); PL – placebo and vitamin D ( $p = 0.9817$ ); and C – vitamin D ( $p = 0.8448$ ). Results represent mean  $\pm$  SD. The results show significant decrease in weight in Group A (glucan and vitamin D), whereas in Group B (placebo and vitamin D) and Group C were seen only insignificant changes

Figure 2 focused on BMI in our groups. The results summarized in Fig. 2 show significant decrease in weight in Group A (glucan and vitamin D), whereas in

Group B (placebo and vitamin D) and Group C were seen only insignificant changes ( $p = 0.9817$  and  $p = 0.8448$ , resp.).

## Discussion

Inflammation and some mechanisms of immune response are two parts participating in development and progression of DR. It has been established that DM significantly affects neurovascular parts of retina in connection with vascular, neural, glial and immune mechanisms (Duh *et al.*, 2017). Detailed elucidation of components mentioned above is based on attempts to find new therapeutic and preventive actions. These components are able to affect individual mechanisms playing a role in retina damage. Potential preventive regenerative steps might better control development of DR including treatment of early stages of the disease. In addition, they could help use better treatment based on individual conditions of each patient. Therefore, detailed evaluation of individual steps leading to developing DR is even more important. It seems elevated weight and particularly its steady increase, represent one of the important risk factors, resulting not only in DM, but also in DR. Significant decrease of glycol hemoglobin, increase of HDL levels and increase of blood pressure are common in obese individuals and represent clear risk factor for DR development (Zhou *et al.*, 2017). Incremental increase of weight is accompanied by an increase of inflammatory proteins levels. Their levels are influenced by increasing numbers of adipocytes and preadipocytes producing numerous inflammatory proteins (Engstrom *et al.*, 2003). A HDS-CC study showed connection among increasing levels of ORM, CRP, haptoglobin and risk of DM development (Muhammad *et al.*, 2017). Inflammatory response results, among others, also in expression of serum amyloid A, is clearly connected with multifunctional apolipoproteins involved in metabolism of cholesterol and modulation of some immune mechanisms (Jensen and Whitehead, 1998). It is assumed that after physical stress, glucose levels are connected to ORM levels even in non-obese individuals. This potential relation is subject of several studies (Tsuboi *et al.*, 2018). Evaluation of ORM levels in DM patients seems to be important most of all due to its wide spectrum of effects in case of DM development (Luo *et al.*, 2015). ORM supports accumulation of glycogen in muscle cells and increases some cellular functions (Qin *et al.*, 2016). ORM regulates food intake and energy homeostasis via leptin receptor signaling pathway (Sun *et al.*, 2016). Based on these findings, one can hypothesize about possible regulation of obesity and/or metabolic problems of ORM expression regulation.

In our experimental group supplemented with glucan and vitamin D, we found significant increase in well-being and strong decrease of fatigue, which corresponds to the possibility of ORM modulation in similar conditions (Lei *et al.*, 2016). ORM is considered to be an important indicator of an association between obesity and periodontitis (Range *et al.*, 2013). Our experimental group could not confirm these associations, because our

group of patients belong to a population with lower economics, education and health awareness. New findings describing two different members of ORM family, ORM1 and ORM2, offer new possibilities on how to better understand the part of ORM that plays in induction of DM and/or DR. ORM2 is considered to be an anti-inflammatory and immunodulatory agent with anti-neutrophil and anti-complement part, able to inhibit cytokine secretion. In addition, it is involved in regulation of angiogenesis and in regulation of capillary permeability (Duh *et al.*, 2017; Hajer *et al.*, 2008; Fang *et al.*, 2015). ORM1 is assumed to play a role in induction of cancer development and is involved in regulation of angiogenesis (Muhammad *et al.*, 2017). We based our study of ORM in DM2 on published findings of relation between ORM and DM1 (El-Beblawy *et al.*, 2016). Our data showed that evaluation of serum levels of ORM and its dynamics related to DM complication is valuable. In addition, we also tried to measure ORM in urine (unpublished results). Our study showed the evaluation of ORM in urine of patients with DM and DR might be important, which agrees with results of other laboratories (Christiansen *et al.*, 2010; KUSTAN *et al.*, 2016; Talks *et al.*, 2018). In all of our patients, we found significant increase of ORM levels when compared to normal population of the same age. Elevated levels of ORM in patient urine with DR might reflect not only risk of urinary tract infection (which could be eliminated by bacteriological evaluation), but another possible complication: risk of damaged glomerular functions.

In the last decade, Europe and probably the whole world, observed a significant deficit of vitamin D in the entire population (Spiro and Buttriss, 2014). Steadily increasing knowledge about the role of vitamin D deficit in appearance and development of clinical progress of several diseases results in increased interests in this subject; not only from the medical and scientific profession, but also from the general population. Attempts to increase the interests of general public in consumption of food with higher content of vitamin D (either naturally or as a supplement) is increasing (Spiro and Buttriss, 2014; Tessari and Lante, 2017; Aoe *et al.*, 2017). Lately, an association between vitamin D deficit and risk of DM2 development is being discussed in the literature (Lu *et al.*, 2018). Our findings show that vitamin D supplementation helps to improve several clinical manifestation of DM2 and its complications (Zavorkova *et al.*, 2018; Christiansen *et al.*, 2010). Risk of DR development in overweight or obese individuals is well documented (Zhou *et al.*, 2017) and further confirmed by our findings. Our group of DR patients revealed that starting levels of vitamin D are approximately 20% lower when compared with patients with DM2. In addition, even 6 month supplementation did not cause an increase up to normal levels. Dynamics of vitamin D

levels in our group was  $14.2 \pm 5.9$  ng/ml at start,  $22.5 \pm 7.5$  ng/ml after three months and  $29.2 \pm 14.2$  ng/ml after six months. The recommended value (30 ng/ml) was reached only in 36.5% of patients. Based on detailed evaluation, we concluded that the reason was inadequate adherence to recommended conditions of vitamin D supplementation.

Effects of glucan in prevention and treatment of obesity and clinical manifestation of DM2 were tested both in experimental and clinical studies (Hajer *et al.*, 2008; Richter *et al.*, 2018a; 2018b; Lu *et al.*, 2018). Clinical trials confirmed that glucan supplementation resulted in a decrease of glycemic index and lipid metabolism, particularly cholesterol levels (Lu *et al.*, 2018; Sima *et al.*, 2018). Findings of changes in visceral obesity after glucan treatment, accompanied with decrease of weight and BMI are important (Wang *et al.*, 2016). The positive effects of glucan in prevention and treatment of metabolic syndrome resulted in food fortification with glucan (Cloetens *et al.*, 2012; McRae, 2018; El Khoury *et al.*, 2012). It seems that food fortification with high fiber content has positive effects in regulation of metabolism. This can be used in prevention and management of DM (Richter *et al.*, 2018a; 2018b). In addition, we found significant BMI decrease after supplementation with glucan and vitamin D. As the groups with vitamin D, but without glucan, showed no differences between start and end values, it is safe to assume that glucan is the supplement responsible for this change. Another important effect of glucan are improvements of salivary immunity (McFarlin *et al.*, 2013) and gastrointestinal tract microbiom with positive effects on risk factors of cardiovascular diseases (Fang *et al.*, 2015).

## Conclusion

This study is a follow-up of previous observation showing effects of glucan on lipid metabolism and obesity in patients with DAR (Richter *et al.*, 2018a; 2018b; Zavorkova *et al.*, 2018). In this study we report significant decrease of ORM levels after the end of the three-month treatment. Our results support the fact that continual supplementation with glucan is necessary for DM and DR patients. We understand that in a population with poor dietary habits, this might be difficult. A direct addition of glucan and fiber into some food could represent a solution to this problem.

## Acknowledgment

The authors are thankful to Ms. Tracey Bender for her editorial help.

## Author's Contributions

All authors equally contributed to this study.

## Ethics

This study was Institutional Review Board (Regional Masaryk University) approved and performed in full agreement with the Helsinki declaration.

## References

- Aoe, S., Y. Ichinose, N. Kohyama, K. Komae and A. Takahashi *et al.*, 2017. Effects of high beta-glucan barley on visceral fat obesity in Japanese individuals: A randomized, double-blind study. *Nutrition*, 42: 1-6. DOI: 10.1016/j.nut.2017.05.002
- Bashir, K.M.I. and J.S. Choi, 2017. Clinical and physiological perspectives of beta-glucans: The past, present and future. *Int. J. Mol. Sci.*, 18: 1906-1906. DOI: 10.3390/ijms18091906
- Cloetens, L., M. Ulmius, A. Johansson-Persson, B. Åkesson and G. Önning, 2012. Role of dietary beta-glucans in the prevention of the metabolic syndrome. *Nutr. Rev.*, 70: 444-458. DOI: 10.1111/j.1753-4887.2012.00494.x
- Christiansen, M.S., E. Hommel, L. Friberg, J. Mølvig and E. Magid *et al.*, 2010. Increased urinary orosomucoid excretion is not related to impaired renal function in patients with type 2 diabetes. *J. Diabetes Complicat.*, 24: 28-36. DOI: 10.1016/j.jdiacom.2008.08.001
- Duh, E.J., J.K. Sun and A.W. Stitt, 2017. Diabetic retinopathy: Current understanding, mechanisms and treatment strategies. *JCI Insight*. 2: e93751-e93751. DOI: 10.1172/jci.insight.93751
- El-Beblawy, N.M., N.G. Andrawes, E.A. Ismail, B. El-Said Enany and H.S.A. El-Seoud *et al.*, 2016. Serum and urinary orosomucoid in young patients with type 1 diabetes: a link between inflammation, microvascular complications and subclinical atherosclerosis. *Clin. Appl. Thromb. Hemost.*, 22: 718-726. DOI: 10.1177/1076029616637185
- El Khoury, D., C. Cuda, B.L. Luhovyy and G.H. Anderson, 2012. Beta glucan: Health benefits in obesity and metabolic syndrome. *J. Nutr. Metab.* DOI: 10.1155/2012/851362
- Engstrom, G., B. Hedblad, L. Stavenow, P. Lind and L. Janzon *et al.* 2003. Inflammation-sensitive plasma proteins are associated with future weight gain. *Diabetes*, 52: 2097-2101. DOI: 10.2337/diabetes.52.8.2097
- Fang, T., M. Cui, J. Sun, C. Ge and F. Zhao *et al.*, 2015. Orosomucoid 2 inhibits tumor metastasis and is upregulated by CCAAT/enhancer binding protein beta in hepatocellular carcinomas. *Oncotarget*, 6: 16106-61619. DOI: 10.18632/oncotarget.3867

- Garcia-Bailo, B., A. El-Sohemy, P.S. Haddad, P. Arora and F. BenZaied *et al.*, 2011. Vitamins D, C and E in the prevention of type 2 diabetes mellitus: modulation of inflammation and oxidative stress. *Biologics*, 5: 7-19. DOI: 10.2147/BTT.S14417
- Garcia-Munoz, A., M.A. Rodriguez, R. Bologna-Molina, F.E. Cázares-Raga and F.C. Hernández-Hernández *et al.*, 2012. The orosomucoid 1 protein (alpha1 acid glycoprotein) is overexpressed in odontogenic myxoma. *Proteome Sci.*, 10: 49-49. DOI: 10.1186/1477-5956-10-49
- Hajer, G.R., T.W. van Haefen and F.L. Visseren, 2008. Adipose tissue dysfunction in obesity, diabetes and vascular diseases. *Eur. Heart J.*, 29: 2959-2971. DOI: 10.1093/eurheartj/ehn387
- Jensen, L.E. and A.S. Whitehead, 1998. Regulation of serum amyloid a protein expression during the acute-phase response. *Biochem. J.*, 334: 489-503. DOI: 10.1042/bj3340489
- Kustan, P., B. Szirmay, Z. Horvath-Szalai, A. Ludány and A. Lakatos *et al.*, 2016. Urinary orosomucoid: Validation of an automated immune turbidimetric test and its possible clinical use. *Biochem. Med. (Zagreb)*, 26: 421-430. DOI: 10.11613/BM.2016.044
- Lei, H., Y. Sun, Z. Luo, G. Yourek and H. Gui *et al.*, 2016. Fatigue-induced orosomucoid 1 acts on C-C chemokine receptor type 5 to enhance muscle endurance. *Sci. Rep.*, 6: 18839-18839. DOI: 10.1038/srep18839
- Lu, L., D.A. Bennett, I.Y. Millwood, S. Parish and M.I. McCarthy *et al.*, 2018. Association of vitamin D with risk of type 2 diabetes: A Mendelian randomisation study in European and Chinese adults. *PLoS Med.*, 15: e1002566-e1002566. DOI: 10.1371/journal.pmed.1002566
- Luo, Z., H. Lei, Y. Sun, X. Liu and D.F. Su, 2015. Orosomucoid, an acute response protein with multiple modulating activities. *J. Physiol. Biochem.*, 71: 329-340. DOI: 10.1007/s13105-015-0389-9
- Lyon, M.R. and R. Reichert, 2010. The effect of a novel viscous polysaccharide along with lifestyle changes on short-term weight loss and associated risk factors in overweight and obese adults: An observational retrospective clinical program analysis. *Altern. Med. Rev.*, 15: 68-75. PMID: 20359270
- McFarlin, B.K., K.C. Carpenter, T. Davidson and M.A. McFarlin, 2013. Baker's yeast beta glucan supplementation increases salivary IgA and decreases cold/flu symptomatic days after intense exercise. *J. Diet. Suppl.*, 10: 171-183. DOI: 10.3109/19390211.2013.820248
- McRae, M.P., 2018. Dietary fiber intake and type 2 diabetes mellitus: An umbrella review of meta-analyses. *J. Chiropr. Med.*, 17: 44-53. DOI: 10.1016/j.jcm.2017.11.002
- Muhammad, I.F., Y. Borne, G. Ostling, C. Kennbäck and M. Gottsäter *et al.*, 2017. Acute phase proteins as prospective risk markers for arterial stiffness: The Malmo Diet and Cancer cohort. *PLoS One*, 12: e0181718-e0181718. DOI: 10.1371/journal.pone.0181718
- Nita, M., A. Grzybowski and F.J. Ascaso, 2014. Age-related macular degeneration in the aspect of chronic low-grade inflammation (pathophysiological parainflammation). *Mediators Inflamm.*, 2014: 930671-930671. DOI: 10.1155/2014/930671
- Qin, Z., J.J. Wan, Y. Sun, P.Y. Wang and D.F. Su *et al.*, 2016. ORM Promotes skeletal muscle glycogen accumulation via CCR5-activated AMPK pathway in mice. *Front. Pharmacol.* DOI: 10.3389/fphar.2016.00302
- Range, H., C. Poitou, A. Boillot, C. Ciangura and S. Katsahian *et al.*, 2013. Orosomucoid, a new biomarker in the association between obesity and periodontitis. *PLoS One*, 8: e57645-e57645. DOI: 10.1371/journal.pone.0057645
- Richter, J., M. Zavorkova, V. Vetvicka, I. Liehneová and V. Kral *et al.*, 2018a. Effects of  $\beta$ -glucan and vitamin D supplementation on inflammatory parameters in patients with diabetic retinopathy. *J. Diet. Suppl.* DOI: 10.1080/19390211.2018.1458769
- Richter, J., M. Závorková, V. Vetvicka, I. Liehneová and V. Král, *et al.*, 2018b. Vitamin D and  $\beta$ -glucan supplementation affects levels of leptin, apolipoproteins and general nutrition state in patients with diabetic retinopathy. *Edorium J. Pathol.* DOI: 10.5348/100009P03RJ2018OA
- Sima, P., L. Vannucci and V. Vetvicka, 2018.  $\beta$ -Glucans and cholesterol (Review). *Int. J. Mol. Med.*, 41: 1799-1808. DOI: 10.3892/ijmm.2018.3411
- Spiro, A. and J.L. Buttriss, 2014. Vitamin D: An overview of vitamin D status and intake in Europe. *Nutr. Bull.*, 39: 322-350. DOI: 10.1111/mbu.12108
- Straczkowski, M., A. Nikolajuk, R. Majewski, R. Filarski and M. Stefanowicz *et al.*, 2018. The effect of moderate weight loss, with or without (1, 3)(1, 6)-beta-glucan addition, on subcutaneous adipose tissue inflammatory gene expression in young subjects with uncomplicated obesity. *Endocrine*, 61: 275-284. DOI: 10.1007/s12020-018-1619-z
- Sun, Y., Y. Yang, Z. Qin, J. Cai and X. Guo *et al.*, 2016. The acute-phase protein orosomucoid regulates food intake and energy homeostasis via leptin receptor signaling pathway. *Diabetes*, 65: 1630-1641. DOI: 10.2337/db15-1193
- Talks, B.J., S.B. Bradwell, J. Delamere, W. Rayner and A. Clarke *et al.*, 2018. Urinary alpha-1-acid glycoprotein is a sensitive marker of glomerular protein leakage at altitude. *High Alt. Med. Biol.*, 19: 295-298. DOI: 10.1089/ham.2018.0017

- Tessari, P. and A. Lante, 2017. A multifunctional bread rich in beta glucans and low in starch improves metabolic control in type 2 diabetes: A controlled trial. *Nutrients*, 9: 297-297.  
DOI: 10.3390/nu9030297
- Tsuboi, A., S. Minato, M. Yano, M. Takeuchi and K. Kitaoka *et al.*, 2018. Association of serum orosomucoid with 30-min plasma glucose and glucose excursion during oral glucose tolerance tests in non-obese young Japanese women. *BMJ Open Diabetes Res. Care*, 6: e000508-e000508.  
DOI: 10.1136/bmjdr-2018-000508
- Wang, Y., N.P. Ames, H.M. Tun, S.M. Tosh and P.J. Jones *et al.*, 2016. High molecular weight barley  $\beta$ -glucan alters gut microbiota toward reduced cardiovascular disease risk. *Front. Microbiol.*, 7: 129-129. DOI: 10.3389/fmicb.2016.00129
- Wolowczuk, I., C. Verwaerde, O. Viltart, A. Delanoye and M. Delacre *et al.*, 2008. Feeding our immune system: Impact on metabolism. *Clin. Dev. Immunol.*, 2008: 639803-639803.  
DOI: 10.1155/2008/639803
- Zavorkova, M., V. Vetvicka, J. Richter, V. Kral and I. Liehnova *et al.*, 2018. Effects of glucan and vitamin D supplementation on obesity and lipid metabolism in diabetic retinopathy. *Open Biochem. J.*, 12: 36-45.  
DOI: 10.2174/1874091X01812010036
- Zhou, Y., Y. Zhang, K. Shi and C. Wang, 2017. Body mass index and risk of diabetic retinopathy: A meta-analysis and systematic review. *Medicine (Baltimore)*, 96: e6754-e6754.  
DOI: 10.1097/MD.0000000000006754